		**************************************	50X1-H
	CLASSIFICATION BECE	R E T/SECURITY INFORMATION	J
	CENTRAL INTELLI	GENCE AGENCY	
	INFORMATIO		
COUNTRY	/ USBR	DATE DISTR. /6 JUN 53	
SUBJECT	Early Chemical Warfare Preparation Gas Masks/Decontamination	NO. OF PAGES 3	
PLACE ACQUIRE			
DATE ACQUIRE			
DATE OF 1			F0V4 I
7010 DOCUMENT OF THE MAITED ASS 704, OF T LATION OF 1TO	Converse derentation appropries for enflowal occupies of the states, private for manife partition for the states of the states o	THIS IS UNEVALUATED INFORMATION	50X1-H
Jaaniaires Sv	LAW. THE ALPHONUTION IF THIS FROM 19 PROMIDITIES.	The state of the s	50X1-H
) 30/1-11
			¥
	War Gases		
			·
		The state of the s	
1.	carried out in glass walled carried	of war gases on tats and wogs were	
	size of the animals in test.	the size of which depended upon the	: ∯ 50X1-⊦
• •	size of the animals in test. They were brought into cylinders or sometimes, as was the	the size of which depended upon the the demonstration laboratory in	 ∯ 50X1-⊦
· ,	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record	the size of which depended upon the the demonstration laboratory in case with hydrocyaniz acid, in rubber to the chambers and the effects were ded in reports.	50X1-H
· .	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals.	the size of which depended upon the the demonstration laboratory in case with hydrocyaniz acid, in rubber to the chambers and the effects were ded in reports.	
2. :	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to be believed through the glass and record hydrocyanic acid killed the animals and phosgers produced characteristic policyting, etc. Animals subjected	the size of which depended upon the the demonstration laboratory in case with hydrocyaniz acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach	
2 . .	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic poisoning, etc. Animals subjected the for the or 48 hours. They were	the size of which depended upon the the demonstration laboratory in case with hydrocyaniz acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the common of sterva-	
2.	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to be a sometime observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic and phosgere produced characteristic policyling, etc. Animals subjected the for the or 48 hours. They were were subject to post sorten examinat	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion.	
2. :	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic noisoning, etc. Animals subjected the for the or 48 hours. They were were subject to post acrtem examinat specific laboratory tests had been weach potential war gas in air, food.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion.	50X1-H
2. 3.	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic poisoning, etc. Animals subjected the for the or 48 hours. They were were subject to post acrtem examinat specific laboratory tests had been weach potential war gas in air, food, laboratory training consisted efforces.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. worked out for the identification of c clothing, drugs, etc. A part of the	
2. ;	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic policying, etc. Animals subjected the for the or 48 hours. They were were subject to post morten examinate Specific laboratory tests had been weach potential var gas in air, food, laboratory training consisted of car samples. Each student taking the conseveral unknowns. He did this by go several unknowns. He did this by go several unknowns.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion.	50X1-⊦ •
3. 8	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic policyling, etc. Animals subjected then for 24 or 48 hours. They were were subject to post morten examinat Specific Isboratory tests had been weach potential war gas in air, food, laboratory training consisted of car samples. Each student taking the conseveral unknowns. He did this by go a prescribed order.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. worked out for the identification of clothing, drugs, etc. A part of the crying out these tests on contaminated purse had to identify the contaminants in bing through identification tests in	50X1-H
3. 4	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to be a sometime observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic noisoning, etc. Animals subjected the for the or 48 hours. They were were subject to post morten examinate Specific laboratory tests had been weach potential war gas in air, food, laboratory training consisted of car samples. Each student taking the conseveral unknowns. He did this by go a prescribed order. Thask, distilling, and then determine	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. worked out for the identification of c clothing, drugs, etc. A part of the crying out these tests on contaminated purse had to identify the contaminants in oung through identification tests in wed placing a piece of cloth in a ting the poison in the distillate.	50X1-⊦ •
2. 3. 4. 5. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6.	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic policy at the for 24 or 48 hours. They were were subject to post sorten examinate Specific Isboratory tests had been weach potential war gas in air, food, laboratory training consisted after care samples. Each student taking the conserved unknowns. He did this by go a prescribed order. the tests on crouning involutions, distilling, and then determing the contributions was protected by impregnational contains was protected by impregnation.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. worked out for the identification of c clothing, drugs, etc. A part of the crying out these tests on contaminated purse had to identify the contaminants in oling through identification tests in ved placing a piece of cloth in a ting the poison in the distillate. on with a secret agent which was never	50X1-H
2	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic policyling, etc. Animals subjected the for the or 48 hours. They were were subject to post morten examinate specific laboratory tests had been we each potential war gas in air, food, laboratory training consisted effications. Each student taking the conseveral unknowns. He did this by go a prescribed order. Thank, distilling, and then determine the tests on crotning involutions was protected by impregnation dentified. Samples or crotning samples of contected, including suits, hats, how	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. worked out for the identification of c clothing, drugs, etc. A part of the rrying out these tests on contaminated purse had to identification tests in ling through identification tests in ved placing a piece of cloth in a ling the poison in the distillate. on with a secret agent which was never of articles of clothing were so outs, gloves, underword.	50X1-F 50X1-F 50X1-F
2	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic policyling, etc. Animals subjected then for the or 48 hours. They were were subject to post sorten examinate specific laboratory tests had been weach potential war gas in air, food, laboratory training consisted efficar samples. Each student taking the conseveral unknowns. He did this by go a prescribed order. Thank, distilling, and then determine clothing was protected by impregnation dentified. Samples of contribution of protected we wool, cotton, rubberired fabric, etc.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. forked out for the identification of clothing, drugs, etc. A part of the trying out these tests on contaminated purse had to identify the contaminants in ling through identification tests in wed placing a piece of cloth in a ling the poison in the distillate. on with a secret agent which was never of articles of clothing were so ots, gloves, underwear, etc. The re made of the usual material such as	50X1-H
2	size of the animals in test. They were brought into cylinders or sometimes, as was the balloons. The gases were admitted to observed through the glass and record hydrocyanic acid killed the animals and phosgere produced characteristic policyling, etc. Animals subjected then for the or 48 hours. They were were subject to post sorten examinate specific laboratory tests had been weach potential war gas in air, food, laboratory training consisted efficar samples. Each student taking the conseveral unknowns. He did this by go a prescribed order. Thank, distilling, and then determine clothing was protected by impregnation dentified. Samples of contribution of protected we wool, cotton, rubberired fabric, etc.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. worked out for the identification of clothing, drugs, etc. A part of the rrying out these tests on contaminated were had to identify the contaminants in ling through identification tests in wed placing a piece of cloth in a ting the poison in the distillate. on with a secret agent which was never f articles of clothing were so ots, gloves, underwear, etc. The	50X1-F 50X1-F 50X1-F
2	size of the animals in test. They were brought into- cylinders or sometimes, as was the balloons. The gases were admitted to balloons. The gases and record the for the or 48 hours. They were were subject to post acrtem examinat specific laboratory tests had been we each potential war gas in air, food, laboratory training consisted of car samples. Each student taking the co several unknowns. He did this by go a prescribed order. the tests on crotning involutions, distilling, and then determin Clothing was protected by impregnation identified. Samples of crotected by impregnation clothing was protected by impregnation clothing the countries of clothing so protected we cool, cotton, rubberired fabric, etc. thich was to be used for the protect.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. forked out for the identification of clothing, drugs, etc. A part of the trying out these tests on contaminated purse had to identify the contaminants in ling through identification tests in wed placing a piece of cloth in a ling the poison in the distillate. on with a secret agent which was never of articles of clothing were so ots, gloves, underwear, etc. The re made of the usual material such as	50X1-F 50X1-F 50X1-F
2	size of the animals in test. They were brought into- cylinders or sometimes, as was the balloons. The gases were admitted to balloons. The gases and record the for the or 48 hours. They were were subject to post acrtem examinat specific laboratory tests had been we each potential war gas in air, food, laboratory training consisted of car samples. Each student taking the co several unknowns. He did this by go a prescribed order. the tests on crotning involutions, distilling, and then determin Clothing was protected by impregnation identified. Samples of crotected by impregnation clothing was protected by impregnation clothing the countries of clothing so protected we cool, cotton, rubberired fabric, etc. thich was to be used for the protect.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. forked out for the identification of clothing, drugs, etc. A part of the trying out these tests on contaminated purse had to identify the contaminants in ling through identification tests in wed placing a piece of cloth in a ling the poison in the distillate. on with a secret agent which was never of articles of clothing were so ots, gloves, underwear, etc. The re made of the usual material such as	50X1-F 50X1-F 50X1-F
2	size of the animals in test. They were brought into- cylinders or sometimes, as was the balloons. The gases were admitted to balloons. The gases and record the for the or 48 hours. They were were subject to post acrtem examinat specific laboratory tests had been we each potential war gas in air, food, laboratory training consisted of car samples. Each student taking the co several unknowns. He did this by go a prescribed order. the tests on crotning involutions, distilling, and then determin Clothing was protected by impregnation identified. Samples of crotected by impregnation clothing was protected by impregnation clothing the countries of clothing so protected we cool, cotton, rubberired fabric, etc. thich was to be used for the protect.	the size of which depended upon the the demonstration laboratory in case with hydrocyanic acid, in rubber to the chambers and the effects were ded in reports. within a few seconds. Mustard gas c skin lesions, lung lesions, stomach to these gases were kept under observa- then killed and the organs affected tion. forked out for the identification of clothing, drugs, etc. A part of the trying out these tests on contaminated purse had to identify the contaminants in ling through identification tests in wed placing a piece of cloth in a ling the poison in the distillate. on with a secret agent which was never of articles of clothing were so ots, gloves, underwear, etc. The re made of the usual material such as	50X1-F 50X1-F 50X1-F

ļ

SECRET/SECURITY INTORNATION

	• 2 -	503/4 11118/
	shown very deep. ulcerated skin lesions on men	50X1-HUM
** e ₁	Whose wieting and district the second	50X1-HUM
	TOUR BUILDER TIME, Rhant harhand fire on the minutes in	
	been been a word, and were then taken out after their injuries has been	50044 111154
	prisoners, or laborers in a poison cas factory on avenue well-	50X1-HUM
	IMAD WOLK WAS Under the direction of Dung (a)	50X1-HUM
		50X1-HUM
	a colonel in the Army at that time. He had a big reputation in the USSR in the war gas field.	50X1-HUM
	One Marke	40000
	Gas Masks	
6	. There were two types of Canisters used with one type of gas mask in the	
		50X1-HUM
	the contents of the canister. The one for the civil population contained line, charcoal, and perhaps other ingredients. The contents of the canister	Sec. 19
	for military were entirely secret.	
7	The gas mask consisted of a without advantage	
•	The gas mask consisted of a rubber piece which covered the entire face and perhaps about three-fourths of the scalp. It was connected to the	
	clumey. There were frequent reports that soldiers engaged in long training marches threw away the masks, but not the canvas carrying bags, because	
	or oner weight diff dwawardhess, the German macks	5024 111184
	were smaller, lighter and less uncomfortable.	50X1-HUM
8.		
	The competition to buelle builts in anti-fit two ining and	
	or one rapid adjustment of the gas marks and then manife the ment	
	going about their normal occupations. Gas maks for children in cribs were available and also masks for horses and dogs.	
_		
9•		
	operations, making rounds of the hospitals, writing reports and eventuing	EOV1 LILIM
	patients. compelled to sleep in gas masks two or three times as a part of the training program.	50X1-HUM
10.		
	All canisters were painted khaki color and no way to distinguish between them except by learning the point of issue.	50X1-HUM
	The state of the Military Mere never strail and the state state state	
	organizations.	
11.	We were compelled to spend some time in gas chambers with the regular	
	military asks and with protective clothing.	50X1-HUM
		SUX I-HUIVI
	Decontamination '	
12.	Decontesination of other amount	
	Decontamination of skin exposed to poison gas was effected by soap and hot water. Afterwards a weak solution of iodine was sometimes painted on the exposed area. For field water and the exposed area.	
	Capoula atta for ileid use, trucke with tanilana and a	
	the troops the troops the troops	50X1-HUM
	and whose showers in training.	
13.	Special trucks on wheels were provided for the decontamination of clothing.	
	The distriction of the contract of the contrac	
	required, formaldehyde was used.	
14.	For the decontamination of rums and utensils, a portable sprayer was	
	large ground areas, large sheet metal drums containing powdered lime or a	

SECRET /SECURITY THEORIGINA